



**US Army Corps  
of Engineers®**

Portland District

USACE PMP Update – 19 June 2019

---

## **MAY UPDATE: PINNIPED ABUNDANCE AND SALMON PREDATION AT BONNEVILLE LOCK AND DAM**

Fisheries Field Unit

U.S. Army Corps of Engineers, Bonneville Lock and Dam, Cascade Locks, OR 97014

Correspondence: [Kyle.S.Tidwell@usace.army.mil](mailto:Kyle.S.Tidwell@usace.army.mil)

This is a monthly status report for the pinniped monitoring program at Bonneville Dam and summarizes the observed fish predation and pinniped abundance at Bonneville Dam from 01 May, 2019 through 31 May, 2019. This is the final update for the 2018 – 2019 season. A final report will be compiled and will be available in the fall.

Previous reports and related PMP information can be found at the link below:

<http://pweb.crohms.org/tmt/documents/FPOM/2010/Task%20Groups/Task%20Group%20Pinnipeds/>

*PLEASE NOTE - All data presented here are preliminary as of the status report date. Predation numbers and abundance estimates are unexpanded and will change as data are proofed and analyzed. Final predation estimate data will be expanded to adjust for hours and days not observed as well as “unknown” prey species consumed for the final report. The final report summarizing the results of the 2019 Pinniped Monitoring Program will be available in the fall of 2019.*



**US Army Corps  
of Engineers®**

Portland District

USACE PMP Update – 19 June 2019

---

## **BACKGROUND**

Concerns regarding an increased number of pinnipeds at Bonneville Dam during the fall and winter and their potential associated impacts on endangered salmonids led to a request by NOAA to monitor the abundance and number of fish killed by these pinnipeds. In response to these concerns, and to fulfill the requirements set forth in the Federal Columbia River Power System Biological Opinion (NMFS 2000, 2008) – which outlines operational criteria for dams to protect ESA-listed fish – the U.S. Army Corps of Engineers, the Fisheries Field Unit initiated a fall and winter pinniped monitoring program to fulfill the Reasonable and Prudent Alternatives defined in the predation management strategy of the Biological Opinion and to provide estimates of pinniped abundance, fish predation, and deterrence strategies. This progress report documents the monitoring of pinniped activities at Bonneville Dam from 08 December, 2018 through 31 January, 2019.

Similar to last year's fall and winter monitoring period, we manipulated previously used spring sampling methods to fit the fall and winter conditions of the Columbia River system. We sampled the priority tailrace (as determined by planned winter outages) and sampled only that tailrace four hours per day in a stratified random fashion whenever the daily abundance counts were greater than 20 pinnipeds (as per study plan provided and approved by NOAA, March 2017). The planned outages for winter maintenance in 2018 are to take place at Power House 2, and as such, Power House 1 tailrace was prioritized for fish predation observations through 31 December, 2018.

The spring sampling period began on 1 January, 2019. As such, both power house tailraces are now being monitored using the same stratified random fashion with the four sampling hours being spread across the two tailraces. Methods and assumptions for observations and estimates of fish predation are captured in the previous year's annual report (Tidwell et al. 2018). These methods consist of visual observation of predation events that are incorporated by fish and pinniped species separately into a probability based estimation calculation to assess the mean level of predation each week. Bootstrap sampling of these estimates provide bounded estimates of predation by week, for each fish species, and by each species of pinniped. Due to the in-season nature of this update and the need to QA/QC data prior to analysis, the estimates provided here are the raw, unadjusted, and un-expanded estimates. Final (bounded) estimates will be provided after the season has completed and the data have been reviewed and analyzed.

Estimates of abundance are made by sampling across the tailrace and at known haul-out sites on Bonneville Project in the early morning hours when animals are most easily observed. Each site is independently counted and aggregated to provide a project wide estimate of pinnipeds each day. Counts are interpolated across weekends and holidays.



**US Army Corps  
of Engineers®**

Portland District

USACE PMP Update – 19 June 2019

---

Thus, the data provided herein, are the daily project wide abundance estimates and the raw number of observed fish killed between 01 May, 2019 and 31 May, 2019. A final report with expanded fish predation estimates will be available after 31 May, 2019.

### **PINNIPED ABUNDANCE**

We present abundance data using the maximum number of individuals counted during a comprehensive tailrace point count and interpolated for days not observed. For inter- and intra-year comparison of abundance estimates, we report average daily abundance with standard deviation as measures of variance.


#### ***Abundance: 01 May, 2019– 31 May, 2019***

Steller Sea Lion (SSL; *Eumetopias jubatus*) have been observed at the dam for 22 days since May 1, 2019, with an average daily abundance estimate of  $14.4 \pm \text{SD } 13.7$  (Table 1). An average of  $1.85 \pm \text{SD } 1.5$  California Sea Lions (CSL; *Zalophus californianus*) were observed during the same period (Figure 1). No harbor seals (*Phoca vitulina*) have been observed since 24 January, 2018.


To date, we have documented 50 SSLs as individuals based on the high point count on 25 April. However, due to the low level of branding for SSL the identification of individuals stands at a count of 12 animals.



### CSL Daily Abundance



### SSL Daily Abundance



\* Averages from 6/1 - 12/31 sporadic between years

Figure 1. Comparison of estimated abundance of each pinniped species at Bonneville Dam between the 10-year running average and the current year.


Table 1. Interpolated daily minimum counts of pinnipeds at Bonneville Dam tailraces between 01 May, 2019 and 31 May, 2019.

Species	$\bar{x} \pm S.D.$	Range	n days = 0
SSL	14.4 ± 13.7	0 - 43	9
CSL	1.9 ± 1.5	0 - 6	5



**FISH PASSAGE & PREDATION**

Salmonid passage for May was below the 10-year average (Figure 2). A total of 600 steelhead (*Oncorhynchus mykiss*) passed in May, which is approximately 30% of the 10-year average. Similarly 49,952 Chinook Salmon crossed Bon during the same period which is approximately 51% of the 10-year average.



*Figure 2. Comparison of the ten year average and current passage of all salmonids over Bonneville Dam between 08 December, 2018 and 31 May, 2019. Data obtained from USACE, FPC – www.FPC.org.*

***Fish Predation: 01 May, 2019– 31 May, 2019***

From 01 May, 2019 to 31 May, 2019, a total 428 Chinook Salmon, seven steelhead, four White Sturgeon (*Acipenser transmontanus*), four Pacific Lamprey (*Entosphenus tridentatus*) and 36 “unknown” predation events by pinnipeds have been documented (Table 2). Upon review and expansion for hours not observed and unknown fish predation events, the estimated number of fish killed will increase.



**US Army Corps  
of Engineers®**

Portland District

USACE PMP Update – 19 June 2019

---

---

*Table 2. Observed fish consumption by both species of pinniped at Bonneville Dam from 01 May, 2019 and 31 May, 2019.*

Species	Chinook	Coho	Steelhead	Sturgeon	Lamprey	Other	Unknown	Total
SSL	376	0	7	4	2	2	34	425
CSL	52	0	0	0	2	0	2	56
Total	428	0	7	4	4	2	36	481

### **DISCUSSION**

Relative to the May monitoring conducted last year, SSL abundance was similar and CSL abundances were significantly lower (i.e. CSL average abundance 2018 = 8.15 vs. 2019 = 1.8). The last SSL was documented on 22 May, and the last CSL was observed on 28 May. Fish predation is also similar to the previous year and peaked during the last month of pinniped presence at BON. Monitoring ceased on 31 May after a three day period of pinniped absence. The USACE FFU will monitor for pinniped presence in the fall and winter and respond with monitoring efforts when there are  $\geq 20$  animals present, as directed by the 2019 interim BiOp.

### **LITERATURE**

NMFS (National Marine Fisheries Service). 2000. Federal Columbia River Power System Biological Opinion.

NMFS (National Marine Fisheries Service). 2008. Federal Columbia River Power System Biological Opinion.

Tidwell, K.S., B.K. van der Leeuw, L.N. Magill, B.A. Carrothers, and R. H. Wertheimer. 2018. Evaluation of pinniped predation on adult salmonids and other fish in the Bonneville Dam tailrace, 2017. U.S. Army Corps of Engineers, Portland District Fisheries Field Unit. Cascade Locks, OR. 54pp.